

Dynamics and Scaling of 2D Polymers in a Dilute Solution

E. Falck,^{1,2} O. Punkkinen,^{1,2} I. Vattulainen,^{1,2} and T. Ala-Nissila^{1,3}

¹Laboratory of Physics, Helsinki University of Technology, P.O. Box 1100, FIN-02015 HUT, Finland

²Helsinki Institute of Physics, Helsinki University of Technology, P.O. Box 1100, FIN-02015 HUT, Finland

³Department of Physics, Box 1843, Brown University, Providence R.I. 02912-1843

(Dated: February 2, 2008)

The breakdown of dynamical scaling for a dilute polymer solution in 2D has been suggested by Shannon and Choy [Phys. Rev. Lett. **79**, 1455 (1997)]. However, we show here both numerically and analytically that dynamical scaling holds when the finite-size dependence of the relevant dynamical quantities is properly taken into account. We carry out large-scale simulations in 2D for a polymer chain in a good solvent with full hydrodynamic interactions to verify dynamical scaling. This is achieved by novel mesoscopic simulation techniques.

PACS numbers: 68.35.Fx, 82.20.Wt, 61.20.Ja

The dynamics of polymer chains has attracted attention for decades already. In 3D, polymer dynamics exhibits rich and complex behavior which depends on the solvent conditions and polymer concentration [1, 2]. The 2D case, however, has attracted much less attention. Recently, it has been realized that it has important applications in the field of colloids and biomolecules. Examples include the 2D diffusion of DNA oligonucleotides confined to interfaces [3] and the lateral diffusion of lipids and proteins along biological interfaces [4] such as cell membranes. Further, the dynamics of polymers in 2D is of major importance in thin films whose thickness is less than the size of the polymer. Wetting, surface adhesion, and flow in confined geometries are examples [5] of this broad and fundamental field.

An important feature of essentially all the 2D diffusion processes in soft matter is that they take place in a solvated environment, which implies that the role of the *hydrodynamic interaction* (HI) cannot be disregarded. The HI originates from interactions mediated by the solvent in the presence of momentum conservation. Hydrodynamics plays a major role in a wide range of applications, and hence the understanding of its impact on the dynamics of polymer systems has received a great deal of attention. In 3D the effects of hydrodynamics are well understood: it is well known that the dynamics of polymers in dilute solution is well described by the Zimm model [2]. In 2D, however, the situation becomes significantly more complicated as will be discussed below.

To understand the dynamics of polymer chains, with or without hydrodynamics, a common technique is to apply the theory of dynamical scaling [2]. Two key quantities here are the radius of gyration R_g and the center-of-mass (CM) diffusion coefficient D of the chain. In the dilute limit, they follow the scaling relations $R_g \sim N^\nu$ and $D \sim N^{-\nu_D}$, with corresponding scaling exponents ν and ν_D , respectively. Another central quantity is the intermediate

scattering function defined as

$$S(\vec{k}, t) = (1/N) \sum_{m,n} \left\langle \exp \left\{ i\vec{k} \cdot [\vec{r}_m(t) - \vec{r}_n(0)] \right\} \right\rangle, \quad (1)$$

where N is the degree of polymerization, $\vec{k}$ is a wave vector, and $\{\vec{r}_n\}$'s are the positions of the monomers. This function should then scale as [2]

$$S(k, t) = k^{-1/\nu} F(tk^x), \quad (2)$$

where x is the *dynamical scaling exponent* related to the other exponents through the relation

$$x = 2 + \nu_D/\nu. \quad (3)$$

This is valid for $k \in (2\pi/R_g, 2\pi/a)$, where a is the size of a monomer. Equations (2) and (3) are the cornerstones of dynamical scaling of polymers.

In the purely dissipative case, the values of the scaling exponents for polymer chains are well understood [2]. In the dilute limit the simple Rouse model gives $\nu = 1/2$ and $\nu_D = 1$. When proper volume exclusion is taken into account, $\nu = 3/4$ in 2D and approximately $3/5$ in 3D, while $\nu_D = 1$ still holds for dilute 3D systems and for *all* polymer concentrations in 2D [6].

However, when the HI is taken into account, the situation becomes dramatically different. While in 3D theory and numerical simulations agree with the prediction of the Zimm equations that $\nu = \nu_D$ (*i.e.* $x = 3$) [7, 8, 9, 10], in 2D the situation is less clear. What has been established both theoretically [1, 2] and computationally [11, 12] is that in good solvent conditions $\nu = 3/4$ in 2D, as in the case of no HI.

The situation with ν_D is more subtle, however. Using lattice-gas simulations Vianney *et al.* [12] found a large positive value of $\nu_D = 0.78 \pm 0.05$. The molecular dynamics (MD) simulations of $S(k, t)$ by Shannon and Choy [11], in turn, gave $x = 2$ which would imply that $\nu_D = 0$, if Eq. (3) holds. However, from their MD data for D *vs.* N they concluded that $\nu_D > 0$ thus contradicting the scaling law. They also solved the Zimm equations

numerically in 2D and verified the result that $x = 2$, but found that now $\nu_D < 0$ [11]. These results prompted the authors of Ref. [11] to suggest that dynamical scaling is *broken* for 2D polymers. Essentially, the very basis of polymer dynamics is being questioned.

In this letter, our objective is to determine the validity of dynamical scaling for 2D polymers. To this end, we first present analytic arguments which show that when finite-size effects are properly taken into account, the scaling of D with respect to N is truly logarithmic, leading to $\nu_D = 0$ and thus to $x = 2$. Following this, we carefully extract the exponents ν , ν_D , and x through extensive mesoscopic simulations of a 2D polymer in a good solvent with the full HI included. Our results verify both that $x = 2$ and the predicted logarithmic scaling of D , and thus we conclude that dynamical scaling *is* obeyed in 2D.

To overcome the significant difficulties in simulating polymers with full hydrodynamic interactions, we employ a novel mesoscopic simulation method introduced recently by Malevanets and Kapral (MK) [13, 14]. The MK method is essentially a hybrid molecular dynamics scheme, where the polymer chain is treated microscopically while the solvent obeys coarse-grained dynamics. In practice this idea is implemented by choosing the monomer-monomer and monomer-solvent interactions as in MD simulations, while the conservative interactions between the solvent particles are absent as in the ideal gas. This description preserves the hydrodynamic modes through so-called collision rules. Further, it allows for a major speedup compared to other simulation techniques such as MD.

To describe the dynamics of the coarse-grained solvent, time is partitioned into segments τ and the simulation box is divided into collision volumes or cells. The effective interactions between the solvent molecules take place at each τ : this is called a collision event. In a collision the velocities of the solvent particles are transformed according to $\vec{v}_i(t + \tau) = \vec{V} + \vec{\omega} \cdot [\vec{v}_i(t) - \vec{V}]$. Here $\vec{v}_i$ is the velocity of the particle i , $\vec{V}$ is the average velocity of all the particles in the cell the particle i belongs to, and $\vec{\omega}$ is a random rotation matrix chosen for that particular cell. It can be shown [13] that this multiparticle collision dynamics conserves the momentum and energy in each collision volume, and thus gives a correct description of the hydrodynamics of the velocity field.

Our model system consists of a polymer chain with N monomers immersed in a 2D coarse-grained solvent. The mass of a solvent particle is set to m , and the monomer mass is $2m$. The monomer-monomer and monomer-solvent interactions are described by a truncated Lennard-Jones (LJ) potential:

$$U_{LJ}(r) = \begin{cases} 4\epsilon [(\sigma/r)^{12} - (\sigma/r)^6] + \epsilon, & r \leq 2^{1/6}\sigma; \\ 0, & r > 2^{1/6}\sigma. \end{cases} \quad (4)$$




FIG. 1: The scaling of $S(k, t)$ for a polymer with $N = 40$ in a simulation box with $L = 120$. Here $k \in [1.0, 2.4]$.

Here σ and ϵ together with m define the LJ unit system, where the unit of time is defined as $\tau_{LJ} = \sigma \sqrt{m/\epsilon}$. In addition to the LJ potential, there is an attractive FENE potential between the nearest-neighbor monomers:

$$U_C(r) = -(aR_0^2/2) \ln(1 - r^2/R_0^2), \quad (5)$$

where $a = 7\epsilon\sigma^{-2}$ and $R_0 = 2\sigma$.

The solvent density was set to $\rho = 0.581\sigma^{-2}$, and the simulations were carried out at a temperature $k_B T = 1.2\epsilon$, yielding good solvent conditions. The equations of motion were integrated using the velocity Verlet algorithm with a time step $\delta t = 0.005\tau_{LJ}$. The choice of the parameters that determine the collision dynamics fixes the properties of the coarse-grained solvent, *e.g.* its viscosity. Here we set the collision time to $\tau = \tau_{LJ}$ and the linear size of the collision volume to $l_c = 2\sigma$. The random rotation angles were chosen from a uniform distribution in $[0, 2\pi)$. The size of the polymer chain N varies from 20 to 80 monomers, and the linear system size L ranges from 40σ up to 420σ . Periodic boundary conditions were employed for all system sizes. The CM diffusion coefficient D was determined using the memory expansion method presented in Ref. [15].

First, we checked the scaling of R_g with $N \in [20, 80]$ [16]. Our estimate for the scaling exponent of the radius of gyration is $\nu = 0.75 \pm 0.02$, in excellent agreement with theory. Next, we computed the dynamic structure factor $S(k, t)$ which is depicted in Fig. 1. Our data show the best collapse with $x = 2.0 \pm 0.1$, and we find that x is not particularly sensitive to either N or L when $N < L$ (data not shown). This confirms the MD results of Shannon and Choy [11], and shows that the 3D Zimm result $x = 3$ is indeed invalid in 2D.

Next, we address the crucial question of the value of the scaling exponent ν_D for the CM diffusion coefficient. In the case of the corresponding system in 3D [8], the finite-

size dependence of D is given by $D \sim 1/L$. Therefore, in principle it is easy to determine D for a fixed chain length N by running a series of simulations for different values of L , and then extrapolating to $L \rightarrow \infty$. By repeating this procedure for several values of N , the exponent ν_D can be determined.

However, in the 2D case the finite-size effects are much more subtle due to the infinite range of the HI. We have carefully calculated D analytically for a 2D polymer in a finite system of size L using various approximations, including the approaches presented in Refs. [8, 11, 17]. In all cases we find [18] that D follows the scaling relation

$$D \sim A \ln N - B \ln \frac{1}{L}, \quad (6)$$

where A and B are constants whose values depend on the approximations used [18, 19]. This shows that extracting ν_D in the “traditional” sense in the thermodynamic limit $L \rightarrow \infty$ is no longer possible.

To numerically study the scaling of D , we determined D for each $N \in [20, 80]$ with different values of L . For instance, for $N = 30$ we considered the cases $L \in \{60, 90, 120, 150, 180, 210, 240\}$. For every N , we examined the behavior of D as a function of $\ln(1/L)$, and found that the behavior indeed is linear. Furthermore, we can estimate the exponent ν_D in terms of effective diffusion coefficients in the following way. We chose cutoff values $L_{cut} \in \{10^2, 10^3, 10^4, 10^5, 10^6\}$, and extrapolated a value $D(N, L_{cut})$ for each chain size and cutoff. If the data complied with Eq. (6), we should, when plotting $D(N, L_{cut})$ vs. $\ln N$, obtain a set of equally spaced straight lines. Each line corresponds to a certain cutoff, and the lines should all have the same slope A . As can be seen in Fig. 2, this indeed holds within the statistical uncertainties of our data.

Most importantly, Fig. 2 confirms the prediction of logarithmic scaling of D with N , which means that $\nu_D = 0$. To quantify this, we can extract the exponent ν_D from $\ln D(N, L_{cut})$ vs. $\ln N$: for large values of L , we should have $\ln D \sim -\nu_D \ln N$. The results in Fig. 2 show that ν_D decreases steadily with L as it should. For the largest L_{cut} studied here, we find $\nu_D \approx 0.05 \pm 0.05$.

The analysis above reveals the reason for the suggested breakdown of scaling in Refs. [11, 12]. While the result $x = 2$ is correct, as verified here, the results in the previous studies for ν_D are simply incorrect because the exponent has been extracted without proper finite-size scaling analysis. Thus, we can conclude that dynamical scaling holds for 2D polymers with $x = 2$, $\nu = 3/4$, and $\nu_D = 0$.

Finally, we wish to discuss the issue of long-time tails in 2D diffusion. The velocity autocorrelation function for a tagged particle i is defined as $\phi(t) = \langle \vec{v}_i(t+t') \cdot \vec{v}_i(t') \rangle$ and its integral $\int_0^\infty dt \phi(t)$ yields D . In the presence of the HI it has been shown that $\phi(t) \sim t^{-1}$ [20], or $\phi(t) \sim [t\sqrt{\ln(t)}]^{-1}$ [21], which means that rigorously speaking,




FIG. 2: The dependence of D on the degree of polymerization N for different cutoffs L_{cut} . The lines correspond to $L_{cut} \in \{10^2, 10^3, 10^4, 10^5, 10^6\}$ from bottom to top.

D is not well defined in 2D. This would seem to invalidate the present scaling arguments. However, there are several ways to resolve this problem. A standard method is to view the diffusion coefficients in 2D as time-dependent quantities $D(t)$ [21], from which one can define effective values of $D \equiv D(t_f)$ at some finite time t_f . Alternatively, one can normalize $D(t)$'s with an appropriately chosen “bare” diffusion coefficient $D_0(t)$ (e.g. that of a monomer [6]). Such normalized values $D(t)/D_0(t)$ should converge to a finite result at long times.

However, there are many cases where it is in fact very difficult to observe long-time tails in either experiments [22] or in simulations. Further complications may arise from the fact that confined geometries can strongly influence the asymptotic decay of $\phi(t)$ [23]. In the present case, the issue of long-time tails is settled by recognizing that in dynamical scaling, the absolute values of the diffusion coefficients are irrelevant: only the behavior of D as a function of the chain or system size matters. Hence, we can here use such effective values provided that they have been determined in a consistent way. To this end, we have simply determined D 's over a time interval where the coefficients have converged within numerical error. More precisely, assuming $D \sim \ell_D^2/t_D$, where ℓ_D is the distance over which the chain diffuses during the time interval t_D , the diffusion coefficients have been measured at a point where the chain has diffused a scaled distance




FIG. 3: The convergence of $D(t)$ vs. time for several systems with different values of N and L . The diffusion coefficients have been computed using the memory expansion method (see Ref. [15] for details).

$\ell_D/R_g = 2 - 4$. The convergence of $D(t)$ on these time scales is demonstrated in Fig. 3 for several systems with different values of N and L .

In conclusion, we have applied the MK algorithm to a dilute 2D polymer solution. The MK method itself has proved to be an efficient tool for studies of macromolecular systems, especially in the dilute limit where the computational cost is mainly due to the explicit solvent. The technique has enabled us—at a moderate computational cost—to study system sizes that have not been previously amenable to simulations. This approach together with proper finite-size scaling analysis has allowed us to solve the controversy regarding the dynamical scaling of dilute polymer solutions in 2D with full hydrodynamics. We have found, in contrast to previous arguments [11, 12], that the exponent relation $x = 2 + \nu_D/\nu$ is valid within numerical error. This justifies the scaling hypothesis, and shows that the anomalous exponent $x = 2$ found in previous studies is due to the logarithmic scaling of D as a function of N .

Acknowledgments – This work has been supported in part by the Academy of Finland through its Center of Excellence Program and the National Graduate School in Materials Physics (E.F.).

-
- [1] P.-G. de Gennes, *Scaling Concepts in Polymer Physics* (Cornell University Press, London, 1979).
 - [2] M. Doi and S. F. Edwards, *The Theory of Polymer Dynamics* (Clarendon Press, Oxford, 1986).
 - [3] B. Maier and J. O. Rädler, Phys. Rev. Lett. **82**, 1911 (1999); B. Maier and J. O. Rädler, Macromolecules **33**, 7185 (2000); B. Maier and J. O. Rädler, Macromolecules **34**, 5723 (2001); V. Chan, D. J. Graves, P. Fortina, and

- S. E. McKenzie, Langmuir **13**, 320 (1997).
- [4] P. Cicuta and I. Hopkinson, J. Chem. Phys. **114**, 8659 (2001); M. J. Saxton and K. Jacobson, Annu. Rev. Biophys. Biomol. Struct. **26**, 373 (1997); I. Vattulainen and O. G. Mouritsen, *Diffusion in Membranes*, in *Diffusion in Condensed Matter*, 2nd edition, edited by J. Kärgner and P. Heitjans (Springer-Verlag, in press); Th. Schmidt, G. J. Schütz, W. Baumgartner, H. J. Gruber, and H. Schindler, Proc. Natl. Acad. Sci. USA **93**, 2926 (1996). A. Sonnleitner, G. J. Schütz, and Th. Schmidt, Biophys. J. **77**, 2638 (1999).
- [5] A. Oron, S. H. Davis, and S. G. Bankoff, Rev. Mod. Phys. **69**, 931 (1997).
- [6] T. Ala-Nissila, S. Herminghaus, T. Hjelt, and P. Leiderer, Phys. Rev. Lett. **76**, 4003 (1996); T. Hjelt, S. Herminghaus, T. Ala-Nissila, and S. C. Ying, Phys. Rev. E **57**, 1864 (1998).
- [7] B. Dünweg and K. Kremer, Phys. Rev. Lett. **66**, 2996 (1991); B. Dünweg, J. Chem. Phys. **99**, 6977 (1993).
- [8] B. Dünweg and K. Kremer, J. Chem. Phys. **99**, 6983 (1993).
- [9] C. Pierleoni and J.-P. Ryckaert, Phys. Rev. Lett. **66**, 2992 (1991); C. Pierleoni and J.-P. Ryckaert, J. Chem. Phys. **96**, 8539 (1992).
- [10] P. Ahrlich and B. Dünweg, J. Chem. Phys. **111**, 8225 (1999).
- [11] S. R. Shannon and T. C. Choy, Phys. Rev. Lett. **79**, 1455 (1997).
- [12] J. M. Vianney and A. Koelman, Phys. Rev. Lett. **64**, 1915 (1990).
- [13] A. Malevanets and R. Kapral, J. Chem. Phys. **110**, 8605 (1999).
- [14] A. Malevanets and R. Kapral, J. Chem. Phys. **112**, 7260 (2000).
- [15] S. C. Ying, I. Vattulainen, J. Merikoski, T. Hjelt, and T. Ala-Nissila, Phys. Rev. B **58**, 2170 (1999).
- [16] While we used the original MK method to determine the exponents ν and x , the simplification proposed by A. Malevanets and J. M. Yeomans [Europhys. Lett. **52**, 231 (2000)], was applied for the largest system sizes to compute D , and consequently ν_D . We tested that the CM diffusion coefficients computed by the two versions of the MK algorithm show quantitative agreement.
- [17] R. Kapral, D. Ng, and S. G. Whittington, J. Chem. Phys. **64**, 539 (1976).
- [18] E. Falck, O. Punkkinen, I. Vattulainen, and T. Ala-Nissila, (unpublished).
- [19] This result should be contrasted to the infinite system size result of Eq. (12) in Ref. [11] which also gives logarithmic dependence on N , but becomes negative with increasing N , which is unphysical.
- [20] B. J. Alder and T. E. Wainwright, Phys. Rev. A **1**, 18 (1970).
- [21] T. E. Wainwright, B. J. Alder, and D. M. Gass, Phys. Rev. A **4**, 233 (1971); C. P. Lowe and D. Frenkel, Physica A **220**, 251 (1998).
- [22] M. A. van der Hoef, D. Frenkel, and A. J. C. Ladd, Phys. Rev. Lett. **67**, 3459 (1991).
- [23] M. H. J. Hagen, I. Pagonabarraga, C. P. Lowe, and D. Frenkel, Phys. Rev. Lett. **78**, 3785 (1997); I. Pagonabarraga, M. H. J. Hagen, C. P. Lowe, and D. Frenkel, Phys. Rev. E **59**, 4458 (1999).